

IFT-P.021/99

A NOTE ON THE SUPERSTRING BRST OPERATOR

José N. Acosta¹,
 Nathan Berkovits²,
 and
 Osvaldo Chandía³

*Instituto de Física Teórica, Universidade Estadual Paulista
 Rua Pamplona 145, 01405-900, São Paulo, SP, Brasil*

February, 1999

Abstract

We write the BRST operator of the N=1 superstring as $Q = e^{-R}(\frac{1}{2\pi i} \oint dz \gamma^2 b)e^R$ where γ and b are super-reparameterization ghosts. This provides a trivial proof that Q is nilpotent.

¹e-mail: jose@ift.unesp.br

²e-mail: nberkovi@ift.unesp.br

³e-mail: chandia@ift.unesp.br

1 Introduction

Superstring theory in ten dimensions is a critical $N = 1$ superconformal theory. It can be quantized using a nilpotent BRST operator

$$Q = \frac{1}{2\pi i} \oint dz [c(T_m + \frac{1}{2}T_g) + \gamma(G_m + \frac{1}{2}G_g)] \quad (1)$$

where $[T_m, G_m]$ are the $c = 15$ $N=1$ superconformal generators and $[T_g, G_g]$ are the $c = -15$ $N=1$ superconformal generators constructed from a pair of fermionic ghosts $[b, c]$ and a pair of bosonic ghosts $[\beta, \gamma]$. Physical states are described by vertex operators in the cohomology of Q and, in order to construct vertex operators for the spacetime fermions, it is convenient to fermionize the bosonic ghosts as

$$\beta = \partial\xi e^{-\phi} \quad (2)$$

$$\gamma = \eta e^{\phi}, \quad (3)$$

where η and ξ are free fermions and ϕ is a chiral boson.[1]

Because the above fermionization involves $\partial\xi$ rather than ξ , it is not possible to write the zero mode of ξ in terms of the $[\beta, \gamma]$ ghosts. The Hilbert space without the ξ zero mode is called the “small” Hilbert space, while the Hilbert space including the ξ zero mode is called the “large” Hilbert space. The small Hilbert space can be defined as operators annihilated by $\oint dz \eta$, and one can show that any such operator can be constructed out of the original $[\beta, \gamma]$ ghosts. Since physical vertex operators should be in the small Hilbert space, they must be annihilated by both Q and $\oint dz \eta$. For consistency, this requires that Q should not only be nilpotent, but should also anti-commute with $\oint dz \eta$.

In this letter, we will construct a similarity transformation R such that $Q = e^{-R}(\frac{1}{2\pi i} \oint dz \gamma^2 b)e^R$. (Note that the first term of R was constructed in [2].) Since $\gamma^2 b$ is nilpotent, this trivially proves that Q is nilpotent. Furthermore, since $\oint dz \gamma^2 b$ has trivial cohomology, it proves that Q has trivial cohomology in the large Hilbert space.

However, R does not commute with $\oint dz \eta$, so Q has non-trivial cohomology in the small Hilbert space as expected. Also, $e^{-R}(\frac{1}{2\pi i} \oint dz \gamma^2 b)e^R$ only anti-commutes with $\oint dz \eta$ in the critical dimension, so one cannot use the nilpotent $e^{-R}(\frac{1}{2\pi i} \oint dz \gamma^2 b)e^R$ to quantize the superstring when $D \neq 10$.

2 Similarity transformation

After fermionizing the $[\beta, \gamma]$ ghosts as in (3) and bosonizing $\xi = e^\chi$ and $\eta = e^{-\chi}$, the BRST charge of (1) can be written as $Q = \frac{1}{2\pi i} \oint dz j_{BRST}$ where

$$\begin{aligned} j_{BRST} = & c[T_m - b\partial c - \partial^2\phi - \frac{1}{2}(\partial\phi)^2 + \frac{1}{2}\partial^2\chi + \frac{1}{2}(\partial\chi)^2] \\ & + e^{\phi-\chi}G_m - be^{2(\phi-\chi)} + \partial^2c + \partial(c\partial\chi). \end{aligned} \quad (4)$$

We will now show that

$$j_{BRST} = e^{-R}j_0e^R \quad (5)$$

where

$$j_0 = -be^{2(\phi-\chi)}, \quad (6)$$

$$R = \frac{1}{2\pi i} \oint dz [cG_me^{-\phi}e^\chi - \frac{1}{4}\partial(e^{-2\phi})e^{2\chi}c\partial c]. \quad (7)$$

Note that j_{BRST} was used in [3] as the fermionic generator G^+ of a twisted N=2 superconformal algebra. Using (5), j_{BRST} is trivially nilpotent since j_0 has no poles with itself.

To prove (5), we use the expansion

$$\begin{aligned} e^{-R}j_0e^R &= \sum_{n=0}^{\infty} \frac{1}{n!} j_n, \\ j_n &= [j_{n-1}, R], \end{aligned} \quad (8)$$

where, for $R = \frac{1}{2\pi i} \oint dz r(z)$, the commutator is computed following the rule

$$[j_{n-1}(y), R] = \frac{1}{2\pi i} \oint dz j_{n-1}(y)r(z). \quad (9)$$

If D is the spacetime dimension of the superstring (i.e. $G_m(y)G_m(z) \rightarrow D(y-z)^{-3} + \dots$), the $n=1$ term of (8) is given by

$$\begin{aligned} j_1 = & e^\phi e^{-\chi}G_m + bc\partial c - \frac{3}{2}\partial^2c + \partial c(5\partial\phi - 4\partial\chi) \\ & + c[\frac{3}{2}\partial^2\phi - 3(\partial\phi)^2 - 2(\partial\chi)^2 - \partial^2\chi + 5\partial\phi\partial\chi], \end{aligned} \quad (10)$$

the $n = 2$ term is given by

$$\begin{aligned}
j_2 &= 2cT_m + \frac{D}{2}[\partial^2 c + 2\partial c(\partial\chi - \partial\phi) \\
&+ c(\partial^2\chi - \partial^2\phi + (\partial\phi)^2 + (\partial\chi)^2 - 2\partial\phi\partial\chi)] \\
&- e^{-\phi}e^\chi G_m c\partial c + \frac{5}{4}e^{-2\phi}e^{2\chi}c\partial c\partial^2 c,
\end{aligned} \tag{11}$$

the $n = 3$ term is given by

$$j_3 = 3e^{-\phi}e^\chi G_m c\partial c - \frac{3D}{4}e^{-2\phi}e^{2\chi}c\partial c\partial^2 c, \tag{12}$$

and the $n = 4$ term is given by

$$j_4 = \frac{3D}{2}e^{-2\phi}e^{2\chi}c\partial c\partial^2 c. \tag{13}$$

The terms for $n > 4$ in the expansion vanish identically since the OPE between j_4 and R has no single poles.

It is straightforward to check that j_{BRST} of (4) is equal to

$$j_0 + j_1 + \frac{1}{2!}j_2 + \frac{1}{3!}j_3 + \frac{1}{4!}j_4 \tag{14}$$

when $D = 10$, so we have proven (5). Note that when $D \neq 10$, the integral of (14) contains the term $(10 - D) \oint dz (\frac{1}{2}c\partial\phi\partial\chi + \frac{1}{16}e^{2(\chi-\phi)}c\partial c\partial^2 c)$. Since this term does not anti-commute with $\oint dz \eta = \oint dz e^{-\chi}$, the integral of (14) can only be used as a BRST charge when $D = 10$ for the reasons stated in the introduction.

3 Acknowledgements

JNA would like to acknowledge financial support from FAPESP grant number 98/12362-2, NB would like to acknowledge partial financial support from CNPq grant number 300256/94-9, and OCH would like to acknowledge financial support from FAPESP grant number 98/02380-3.

References

- [1] D. Friedan, E. Martinec and S. Shenker, *Nucl. Phys* **B271**, 93 (1986).
- [2] N. Berkovits, *Phys. Rev. Lett.* **77**, 2891 (1996), hep-th/9604121.
- [3] N. Berkovits, *Nucl. Phys.* **B420**, 332 (1994), hep-th/9308129.